

Geometric scaling in inclusive eA reactions and nonlinear perturbative QCD

A. Freund,¹ K. Rummukainen,² H. Weigert,¹ and A. Schäfer¹

¹*Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany*

²*NORDITA, DK-2100 Copenhagen, Denmark & Department of Physics,*

P.O.Box 64, FIN-00014 University of Helsinki, Finland

(Dated: February 1, 2008)

In this note we report on geometric scaling in inclusive eA scattering data from the NMC and E665 experiments. We show that this scaling, as well as nuclear shadowing, is expected in the framework of nonlinear pQCD at small x based on a simple rescaling argument for ep scattering.

INTRODUCTION

Perturbative QCD, in particular where based on factorization theorems for high energy processes, has been very successful in describing data from high energy experiments. However, perturbative evolution has almost always been *linear* in the parton densities, which was, and very often still is, quite sufficient to describe high energy scattering.

There are, however, situations where this is bound to fail. Among those are high energy scatterings at energies large enough to create a gluon medium in which multi-gluon correlations can no longer be neglected. This has first been discussed in the context of deep inelastic scattering of electrons off (large) nuclei, where both the large number of nucleons and the high energy lead to an increase in the number of gluons involved in the scattering [1, 2, 3, 4, 5, 6, 7, 8, 9]. In this regime nonlinear effects, leading to a taming or saturation/unitarization of F_2^A , should be included.

In the guise of a very successful phenomenological fit, this idea has even been applied to the ep experiment at HERA with its strong and, if untamed, unitarity violating rise of the proton structure function $F_2^p(x_{bj}, Q^2)$ with decreasing x_{bj} [10, 11]. This has led to the observation of “geometric scaling” satisfied by HERA data in the small x_{bj} region [12]. Our main focus here will be an attempt to extend this geometric scaling idea to nuclei and to check if it is already confirmed by available eA data.

The question of the onset of this saturation/unitarization in inclusive QCD observables with increasing energy (decreasing x_{bj}), in particular in the proton structure function $F_2^p(x_{bj}, Q^2)$, has been the subject of active discussions for many years (see a detailed discussion of this subject in [7, 12] and references therein.). Phenomenologically, the key feature at small x_{bj} is the simple structure of the total γ^*p cross section or $F_2^{p,A}$ that arises directly from the underlying physics picture. Viewed in the infinite momentum frame of the target, the virtual photon splits into a $q\bar{q}$ color-dipole (treated as eikonalized Wilson-lines) with a distribution characterized by Q^2 . This dipole then punches through, and interacts with the target, which it sees as a pancake of infinitesimal longitudinal thickness. Consequently the

cross section appears as a convolution of the square of a photon wave function, which gives the probability to create the $q\bar{q}$ pair, and a dipole cross section, which contains all the information about the target and the strong interaction physics.

The latter carries x_{bj} dependence which arises from gluonic fluctuations which induce an increase in the number of gluons [19]. From this point of view, the target will display a growing density of gluons per transverse area (integrated over the longitudinal extent of the target) that drives the system into a saturation regime. The corresponding formula for the total cross section in γ^*p is then

$$\sigma_{\text{tot}}(x_{bj}, Q^2) = \int d^2z \int_0^1 d\alpha |\psi_{\gamma^*}(\alpha, z^2, Q^2)|^2 \times \sigma_{\text{dipole}}(x_{bj}, z^2) \quad (1)$$

where α is the longitudinal momentum fraction of the q or $\bar{q}$, and z their relative transverse separation. Note that the Q^2 and x_{bj} -dependence is clearly separated into wave function and dipole cross section. This reflects the natural distinction between the transverse and longitudinal directions seen at high energies. Q^2 sets the scale for transverse resolution, but longitudinal information about the target is subsumed in the x_{bj} dependence of the dipole cross section. The idea is that the dipole cross section depends on x_{bj} only via a scale $Q_s(x_{bj})$ as $\sigma_{\text{dipole}}(x_{bj}, z^2) = \sigma_{\text{dipole}}(z^2 Q_s^2(x_{bj}))$. Q_s is the saturation scale, i. e. the scale below which the gluonic content of the target becomes “black” or σ_{dipole} reaches its large size asymptotics.

Together with Eq. (1) this implies that σ_{tot} and hence F_2^p scales according to

$$\sigma_{\text{tot}}(x_{bj}, Q^2) = \sigma_{\text{tot}}\left(x_0, \frac{Q^2}{Q_s^2(x_{bj})} Q_0^2\right). \quad (2)$$

This phenomenon is called geometric scaling and, as already mentioned, was first observed to be present in HERA data [12]. The fit used in [12] assumes an “eikonal” shape for σ_{dipole} that saturates for large dipoles. As x_{bj} decreases this saturation affects smaller and smaller distances, inducing Q_s to grow, a fact which

was parametrized by

$$Q_s(x_{bj}) := \left(\frac{x_0}{x_{bj}}\right)^\lambda Q_0. \quad (3)$$

The theoretical basis for such saturation/unitarization features, with a scaling behavior as in Eq. (2), together with precisely the functional dependence of Q_s as in Eq. (3), lies in nonlinear evolution equations that try to predict this x_{bj} dependence from first principles.

The most promising approach to properly include saturation/unitarization is the JIMWLK equation [7] which resums the leading $\ln(1/x_{bj})$ terms in *all* N-point correlation functions of the participating fields not just the leading correlators as in $k_\perp$ -factorization. This leads to a non-linear renormalization group equation (RGE) for the generating functional for these N-point correlators which slows down the rapid growth of structure functions at small x_{bj} while at the same time preventing the system from drifting into the infrared. This is the basis for a theoretically sound, selfconsistent treatment. Linearizing the JIMWLK equation, one recovers the well-known BFKL equation (see [7, 13] for details) as its low density limit. We thus learn to interpret the loss of infrared safety in BFKL as being due to the absence of any non-linear corrections and the predicted too rapid a growth in $F_2^p(x_{bj}, Q^2)$ as signals of BFKL having exceeded its range of validity.

The full content of the JIMWLK equation can be accessed numerically: being a functional equation of Fokker-Plank type it can be rewritten as a Langevin equation with white noise. This then can be discretized and solved using lattice methods [14]. This yields the x_{bj} dependence of σ_{dipole} and in particular the evolution “rate” λ of Eq. (3).

Of particular interest to us in this context is the fact that the evolution equation is *target independent*. Target dependence arises only from the initial conditions for the x_{bj} dependence of the dipole cross section (related to the initial distribution of gluons in the transverse plane), which will, in particular, be A dependent. These facts will allow us to compare ep and eA scattering and investigate whether one of the basic predictions of this approach, geometric scaling, can also be found in inclusive eA processes.

GEOMETRIC SCALING AND NUCLEAR SHADOWING WITHIN NONLINEAR PQCD.

Before talking about geometric scaling in eA scattering, one should address nuclear shadowing or the suppression of $\frac{1}{A}F_2^A$ to F_2^p at small x_{bj} . If this is at all possible it increases our phenomenological leverage considerably by allowing us to compare experiments with different A and x_{bj} directly. This is of particular importance as the

available eA data do not offer any leverarm in x_{bj} for a given Q^2 or vice versa.

In fact, the leading effect of nuclear shadowing can be easily incorporated in this approach by realizing that the change in transverse size simply changes the overall normalization of σ_{dipole} while the change in longitudinal extent can be incorporated by increasing our initial Q_s by a factor A^δ . In the simplest case, assuming the distribution of spectator partons in the target to be homogeneous, $\delta = 1/3$. Eq. (2) for $F_2 \sim \sigma \cdot Q^2$ then turns into

$$\left(\frac{x_{bj}}{x_0}\right)^{2\lambda} \frac{F_2^A(x_{bj}, Q^2)}{A^{2/3+1/3}} = F_2^p \left(x_0, \left(\frac{x_{bj}}{x_0}\right)^{2\lambda} \frac{Q^2}{A^{1/3}}\right). \quad (4)$$

Note that $0 \leq x_{bj} \leq A$ since x_{bj} is given per nucleon. The power $2/3$ in the overall A dependence stems from a simple geometric factor, the area of the target, in the impact parameter integral in σ_{dipole} . Thus we have an unambiguous prediction for $\frac{1}{A}F_2^A$ which states that not only can the observed nuclear shadowing in the eA data be explained by a simple rescaling of the variable Q^2 with $A^{1/3}$, but also that all $\frac{1}{A}F_2^A$ data should, plotted vs.

$\tau = \left(\frac{x_{bj}}{x_0}\right)^{2\lambda} \frac{Q^2}{A^{1/3}}$, lie on the same curve as the data for F_2^p . Hence, this approach also predicts geometric scaling in inclusive eA scattering analogous to the one observed in ep scattering.

GEOMETRIC SCALING IN $\frac{1}{A}F_2^A$

Note that Eq. (4) is strictly true only in the asymptotic region of small x_{bj} , large Q^2 and large nuclei. Therefore, we expect that there might be nonperturbative corrections which can spoil the form of Eq. (4) and consequently we perform a fit to the combined nuclear data with the following form:

$$\left(\frac{x_{bj}}{x_0}\right)^{2\lambda} \frac{1}{A^\gamma} \frac{1}{A} F_2^A(x_{bj}, Q^2) = F_2^p \left(x_0, \left(\frac{x_{bj}}{x_0}\right)^{2\lambda} \frac{Q^2}{A^\delta}\right), \quad (5)$$

where γ and δ are the fit parameters, which ideally would be 0 and $1/3$ respectively.

The starting point of our analysis is the published NMC [15] and E665 [16, 17] data for the ratios of nuclear to deuterium structure functions $2F_2^A/AF_2^D$. In Fig. 1 we show the ratios as functions of x . From these ratios we obtain the nuclear structure functions F_2^A as follows: we first note that NMC data show the deuterium to proton structure function ratio being $F_2^D/2F_2^p = 1$ to better than 95% accuracy in the dynamical range of interest ($x_{bj} < 0.1$) [15], which is well within the error margins of the nuclear F_2 measurements. Thus, we convert F_2^A/F_2^D to structure functions F_2^A by multiplying the ratios by F_2^p , using the geometric scaling ansatz to extrapolate F_2^p to the appropriate values of x_{bj} and Q^2 .




FIG. 1: Ratio of $2F_2^A/AF_2^D$ vs. τ for NMC and E665. Note the increasing steepness with A within the NMC and E665 datasets respectively.




FIG. 2: Comparative phase space of HERA, NMC and E665.

This rescaling is necessary since the kinematic ranges of the HERA, NMC and E665 experiments are not identical as shown in Fig. 2.

Now we are ready to use Eq. (5) to compare nuclear and proton structure functions. In Fig. 1 the rescaled F_2^A is seen to fall on top of the dashed line representing the geometric scaling fit to the HERA F_2^p . For comparison we also show the quality of the scaling fit within the HERA




FIG. 3: Scaling behavior of NMC and E665 F_2^A data vs. $\tau = \left(\frac{x_{b1}}{x_0}\right)^{2\lambda} \frac{Q^2}{A^{1/3}}$. The vertical axis corresponds to the l.h.s. of Eq. (5). The dashed line corresponds to the geometric scaling curve obtained from HERA data. These are shown offset by a factor of 5.

data, offset by a factor of 5.

The best fit to the combined nuclear data is achieved for $\gamma = 0.09$ and $\delta = 1/4$ which are close to but not quite the asymptotic values the nonlinear pQCD approach predicts. The remaining parameter $\lambda \sim 0.18$ in Eq. (5) is already determined from the geometric scaling fit to the HERA data. With this value for λ , the saturation scale at the upper end of the HERA shadowing region at $x_0 = 10^{-1}$ comes out to be marginally “perturbative”, $Q_s^2 = 1 \text{ GeV}^2$.

We found that the NMC calcium data allows a smaller value of γ and, more importantly, a value of $\delta = 1/3$ if the data on lighter nuclei is left out. This can be seen best in a log-linear plot of the scaled F_2 ratios shown in Fig. 4, where the plateau from $\tau = 0.1$ to $\tau = 10$ shows good overall scaling behavior. $\delta = 1/3$ would lift the NMC Ca data fully up to 1. This seems to indicate that the Ca nucleus is large enough for the basic assumptions of the nonlinear pQCD approach to be valid. A very encouraging sign, indeed.

Also note that there is a consistent difference in the E665 and NMC data with the E665 data being much flatter in the ratio $2F_2^A/AF_2^D$, as evidenced in the comparison between the two calcium data sets and shown in Fig. 4. This apparent inconsistency between the two data sets disappears when the ratio $12F_2^A/AF_2^C$ is compared between the two experiments [18]. This question needs to be addressed experimentally.




FIG. 4: Plot of scaled NMC and E665 F_2^A/AF_2^P data vs. $\tau = \left(\frac{x_{bj}}{x_0}\right)^{2\lambda} \frac{Q^2}{A^{1/3}}$. This corresponds to “dividing the data shown in Fig. 3 by the dashed line.”

Notwithstanding all of the above, nuclear data from the proposed Electron Ion Collider at BNL on heavy nuclei at small x_{bj} and large Q^2 is necessary to be absolutely definitive about geometric scaling in inclusive eA scattering. These experiments will by design provide data at smaller x_{bj} but one should take care to open up the phase space region. Only if we have access to a range of x_{bj} values at any given Q^2 (and vice versa) for each species A , can we obtain independent fits for the evolution rate λ . This would be a prerequisite to experimentally disentangle and confirm the A and x dependence extracted simultaneously above. As the errors on the nuclear data get smaller one would also require a direct measurement of F_2^A/AF_2^P that would allow us to do away with our rather cavalier treatment of $F_2^D/2F_2^P$.

CONCLUSIONS

To summarize, we demonstrate for the first time that the NMC and E665 data for the nuclear structure function $\frac{1}{A}F_2^A$ also exhibit, modulo the caveats given above, geometric scaling as already seen in the nucleon structure function F_2^p . This has been confirmed using the shadowing features of the nonlinear pQCD approach, which, as

the dominant effect, consists of a simple rescaling of Q_s^2 by $A^{1/3}$. With the presently available nuclear data which are at rather low Q^2 we had to allow slight modifications in the powers entering the A dependence in order to get a perfect fit. The NMC data for calcium, the heaviest nucleus available within this dataset, would in fact be compatible with the $A^{1/3}$ scaling. Given the limited phase space regions covered in the data sets we find it very encouraging how well the geometric scaling in eA follows the one in ep .

AF was supported by the Emmy-Noether Grant of the DFG, AS by BMBF and HW by the DFG Habilitanden-program.

-
- [1] L. D. McLerran and R. Venugopalan, Phys. Rev. **D49**, 2233 (1994), hep-ph/9309289.
 - [2] L. D. McLerran and R. Venugopalan, Phys. Rev. **D49**, 3352 (1994), hep-ph/9311205.
 - [3] J. Jalilian-Marian, A. Kovner, L. D. McLerran, and H. Weigert, Phys. Rev. **D55**, 5414 (1997), hep-ph/9606337.
 - [4] I. Balitsky, Nucl. Phys. **B463**, 99 (1996), hep-ph/9509348.
 - [5] J. Jalilian-Marian, A. Kovner, A. Leonidov, and H. Weigert, Phys. Rev. **D59**, 014014 (1999), hep-ph/9706377.
 - [6] J. Jalilian-Marian, A. Kovner, and H. Weigert, Phys. Rev. **D59**, 014015 (1999), hep-ph/9709432.
 - [7] H. Weigert, Nucl. Phys. **A703**, 823 (2002), hep-ph/0004044.
 - [8] E. Iancu, A. Leonidov, and L. D. McLerran, Nucl. Phys. **A692**, 583 (2001), hep-ph/0011241.
 - [9] E. Ferreira, E. Iancu, A. Leonidov, and L. McLerran (2001), hep-ph/0109115.
 - [10] I. Abt et al. (H1), Nucl. Phys. **B407**, 515 (1993).
 - [11] M. Derrick et al. (ZEUS), Phys. Lett. **B316**, 412 (1993).
 - [12] A. M. Stasto, K. Golec-Biernat, and J. Kwiecinski, Phys. Rev. Lett. **86**, 596 (2001), hep-ph/0007192.
 - [13] J. Jalilian-Marian, A. Kovner, A. Leonidov, and H. Weigert, Nucl. Phys. **B504**, 415 (1997), hep-ph/9701284.
 - [14] K. Rummukainen and H. Weigert, in preparation (2002).
 - [15] M. Arneodo et al. (New Muon), Nucl. Phys. **B487**, 3 (1997), hep-ex/9611022.
 - [16] M. R. Adams et al. (E665), Z. Phys. **C65**, 225 (1995).
 - [17] M. R. Adams et al. (E665), Z. Phys. **C67**, 403 (1995), hep-ex/9505006.
 - [18] NMC Collab., private communication.
 - [19] This clean separation is in fact a scheme choice: Depending on the frame these fluctuations may be assigned to the photonic probe or the hadronic target. Within this picture we attribute all of them to the target.